



**Figure 2** Contrast Left Ventriculography in the Right Anterior Oblique Projection

The left ventriculogram at end-diastole (A) and end-systole (B), showing apical ballooning and basal hyperkinesis.

## REFERENCES

1. Bybee KA, Murphy J, Prasad A, et al. Acute impairment of regional myocardial glucose uptake in the apical ballooning (takotsubo) syndrome. *J Nucl Cardiol* 2006;13:244-50.
2. Scholte A, Bax JJ, Stokkel MP, et al. Multimodality imaging to diagnose takotsubo cardiomyopathy. *J Nucl Cardiol* 2006;13:123-6.
3. Kume T, Akasaka T, Kawamoto T, et al. Assessment of coronary microcirculation in patients with takotsubo-like left ventricular dysfunction. *Circ J* 2005;69:934-9.
4. Ito K, Sugihara H, Kinoshita N, et al. Assessment of takotsubo cardiomyopathy (transient left ventricular apical ballooning) using <sup>99m</sup>Tc-tetrofosmin, <sup>123</sup>I-BMIPP, <sup>123</sup>I-MIBG and <sup>99m</sup>Tc-PYP myocardial SPECT. *Ann Nucl Med* 2005;19:435-45.
5. Kurisu S, Inoue I, Kawagoe T, et al. Myocardial perfusion and fatty acid metabolism in patients with takotsubo-like left ventricular dysfunction. *J Am Coll Cardiol* 2003;41:743-8.
6. Abe Y, Kondo M, Matsuoka R, et al. Assessment of clinical features in transient left ventricular apical ballooning. *J Am Coll Cardiol* 2003;41:737-42.

## Letters to the Editor

# Cardiovascular Mortality in High-Risk Patients and T-786C Polymorphism in Promoter Region of the Endothelial Nitric Oxide Synthase Gene

Rossi et al. (1) reported very interesting and intriguing results of the first prospective study examining the possible effects of 2 single nucleotide polymorphisms (SNP) in the endothelial nitric oxide synthase (eNOS) gene (the T-786C SNP in the promoter region, and the G894T SNP in exon 7) on cardiovascular mortality among high-risk patients. Although no significant effects were found for the G894T SNP, more cardiovascular deaths were found when individuals with TT genotype for the T-786C SNP were compared with CC + CT individuals (1). The significant effect of T-786C SNP on cardiovascular mortality persisted even after many confounding factors were taken into consideration. How-

ever, a significant number of individuals (32%) were on lipid-lowering therapy at recruitment (1), and it is probable that an increased proportion of these subjects may have received statins thereafter.

Interestingly, although the T-786C SNP apparently does not significantly affect nitric oxide (NO) availability (2,3), it may modulate the responses to statins. In this regard, we have recently reported that treatment with atorvastatin significantly increased NO availability (measured as whole blood nitrite) in CC individuals, but not in TT individuals (4), thus confirming previous findings suggesting that statins may produce stronger effects on NO availability in CC individuals compared with TT individuals (5). In addition, atorvastatin significantly reduced the concentrations of inflammatory markers in subjects with CC (but not TT) genotype (6). Although these findings derive from studies that included healthy individuals, they suggest that statins may significantly modify the cardiovascular risk associated with the T-786C SNP. Indeed, it is possible that treatment with statins counteracts the effects associated with the T-786C SNP (4), thus leading to apparently paradoxical results such as those reported by Rossi et al. (1).

**\*Jose Eduardo Tanus-Santos, MD, PhD**  
**Antonio Casella-Filho, MD, MSc**

\*Department of Pharmacology  
Faculty of Medicine of Ribeirao Preto  
University of São Paulo  
Av. Bandeirantes, 3900  
14049-900 Ribeirao Preto, SP  
Brazil  
E-mail: tanus@fmrp.usp.br

doi:10.1016/j.jacc.2006.12.030

## REFERENCES

1. Rossi GP, Maiolino G, Zanchetta M, et al. The T(-786)C endothelial nitric oxide synthase genotype predicts cardiovascular mortality in high-risk patients. *J Am Coll Cardiol* 2006;48:1166-74.
2. Nagassaki S, Metzger IF, Souza-Costa DC, Marroni AS, Uzuelli JA, Tanus-Santos JE. eNOS genotype is without effect on circulating nitrite/nitrate level in healthy male population. *Thromb Res* 2005;115:375-9.
3. Metzger IF, Souza-Costa DC, Marroni AS, et al. Endothelial nitric oxide synthase gene haplotypes associated with circulating concentrations of nitric oxide products in healthy men. *Pharmacogenet Genomics* 2005;15:565-70.
4. Nagassaki S, Sertorio JTC, Metzger IF, Bern AF, Rocha JBT, Tanus-Santos JE. eNOS gene T-786C polymorphism modulates atorvastatin-induced increase in blood nitrite. *Free Radic Biol Med* 2006;41:1044-9.
5. Abe K, Nakayama M, Yoshimura M, et al. Increase in the transcriptional activity of the endothelial nitric oxide synthase gene with fluvastatin: a relation with the -786T>C polymorphism. *Pharmacogenet Genomics* 2005;15:329-36.
6. Souza-Costa DC, Sandrim VC, Lopes LF, Gerlach RF, Rego EM, Tanus-Santos JE. Anti-inflammatory effects of atorvastatin: modulation by the T-786C polymorphism in the endothelial nitric oxide synthase gene. *Atherosclerosis* 2007. In press.

# Effects of Endothelial Nitric Oxide Synthase Gene Polymorphisms on Oxidative Stress, Inflammatory Status, and Coronary Atherosclerosis: An Example of a Transient Phenotype

In their recent study, Rossi et al. (1) showed that the T786C polymorphism on the promoter region of the endothelial nitric oxide synthase (eNOS) gene modifies redox-sensitive inflammatory pathways, and may be a predictor for clinical outcome in high-risk patients with coronary atherosclerosis. Although this polymorphism was found to be in linkage disequilibrium with the functional polymorphism G894T ( $D' = 0.3$ ), the latter had no predictive value in this cohort, despite the results of a large meta-analysis suggesting the opposite (2). However, the observation that the 894T/786T haplotype leads to a worse cardiovascular death-free survival supports the hypothesis of a more complex association between these polymorphisms and eNOS function.

We have recently shown (3) that the presence of the 894T allele is associated with higher levels of oxidized low-density lipoprotein

and proinflammatory cytokines only under conditions of "biological stress," such as during the acute phase of myocardial infarction, an effect not observed in the same subjects 1 year after the event, or in healthy individuals. Moreover, the 894T allele seems to be associated with impaired endothelial function in high-risk patients (4) and in healthy smokers (5), but not in healthy, low-risk individuals (5).

In the present study, Rossi et al. (1) actually introduce the hypothesis that the previously observed transient effect of the 894T allele on oxidative stress, inflammatory process, and endothelial function (3-5) could actually be due to its linkage disequilibrium with the 786T allele (which has been suggested to modify eNOS expression). However, this could be due to the complex interaction of both polymorphisms constituting the 894T/786T haplotype. The combination of low eNOS expression (induced by the 786T allele) and increased susceptibility of eNOS to proteolytic cleavage (induced by the 894T allele) (6) could lead to a combined effect on the associated phenotype of nitric oxide bioavailability, and the subsequent alterations of oxidative stress and inflammatory status. Therefore, further molecular studies are required to explore the transient behavior of eNOS genotypes/haplotypes, leading to a different phenotype, depending on the underlying disease state.

**\*Charalambos Antoniades, MD**  
**Dimitris Tousoulis, MD, PhD, FACC**  
**Christodoulos Stefanadis, MD, FACC, FESC**

\*Athens University Medical School  
Hippokraton Hospital  
Vasilissis Sophias 114  
115 28 Athens  
Greece  
E-mail: antoniades@panafonet.gr

doi:10.1016/j.jacc.2006.12.029

## REFERENCES

1. Rossi GP, Maiolino G, Zanchetta M, et al. The T(-786)C endothelial nitric oxide synthase genotype predicts cardiovascular mortality in high-risk patients. *J Am Coll Cardiol* 2006;48:1166-74.
2. Casas JP, Bautista LE, Humphries SE, Hingorani AD. Endothelial nitric oxide synthase genotype and ischemic heart disease: meta-analysis of 26 studies involving 23,028 subjects. *Circulation* 2004;109:1359-65.
3. Antoniades C, Tousoulis D, Vasiliadou C, et al. Genetic polymorphism on endothelial nitric oxide synthase affects endothelial activation and inflammatory response during the acute phase of myocardial infarction. *J Am Coll Cardiol* 2005;46:1101-9.
4. Antoniades C, Tousoulis D, Vasiliadou C, et al. Genetic polymorphism G894T on the eNOS gene is associated with endothelial function and vWF levels in premature myocardial infarction survivors. *Int J Cardiol* 2006;107:95-100.
5. Leeson CP, Hingorani AD, Mullen MJ, et al. Glu298Asp endothelial nitric oxide synthase gene polymorphism interacts with environmental and dietary factors to influence endothelial function. *Circ Res* 2002;90:1153-8.
6. Tesaro M, Thompson WC, Rogliani P, Qi L, Chaudhary PP, Moss J. Intracellular processing of endothelial nitric oxide synthase isoforms associated with differences in severity of cardiopulmonary diseases: cleavage of proteins with aspartate vs. glutamate at position 298. *Proc Natl Acad Sci U S A* 2000;97:2832-5.

## Reply

To explain our (1) intriguing results on the effects of the T-786C (in the promoter) and the G894T (in exon 7) single nucleotide polymorphisms (SNP) of the endothelial nitric oxide (NO) synthase (eNOS)